



Package leaflet: Information for the user

AZOPT® 10 mg/ml eye drops, suspension

Brinzolamide

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or your pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What AZOPT is and what it is used for
2. What you need to know before you use AZOPT
3. How to use AZOPT
4. Possible side effects
5. How to Store AZOPT
6. Contents of the pack and other information

1 What AZOPT® is and what it is used for

AZOPT contains brinzolamide which belongs to a group of medicines called carbonic anhydrase inhibitors. It reduces pressure within the eye.

AZOPT eye drops are used to treat high pressure in the eye. This pressure can lead to an illness called glaucoma.

If the pressure in the eye is too high, it can damage your sight.

2 What you need to know before you use AZOPT

Do not use AZOPT

- if you have severe kidney problems.
- if you are allergic to any of the ingredients of AZOPT. For a full list of ingredients please see section 6.
- if you are allergic to medicines called sulphonamides. Examples include medicines used to treat diabetes and infections and also diuretics (water tablets). AZOPT may cause the same allergy.
- if you have too much acidity in your blood (a condition called hyperchloraemic acidosis).

If you have further questions, ask your doctor for advice.

Warnings and precautions

Talk to your doctor or pharmacist before using AZOPT:

- if you have kidney or liver problems.
- if you have dry eyes or cornea problems.
- if you are taking other sulphonamide medicines

Children and adolescents

AZOPT is not to be used by infants, children or adolescents under 18 years of age unless advised by your doctor.

Other medicines and AZOPT

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

If you are taking another carbonic anhydrase inhibitor (acetazolamide or dorzolamide, see section 1 What AZOPT is and what it is used for), talk to your doctor.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

Women who may become pregnant are advised to use effective contraception during AZOPT treatment. The use of AZOPT is not recommended during pregnancy or breast-feeding. Do not use AZOPT unless clearly indicated by your doctor.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Do not drive or use machines until your vision is clear. You may find that your vision is blurred for a time just after using AZOPT.

AZOPT may impair the ability to perform tasks requiring mental alertness and/or physical coordination. If affected, take care when driving or using machines.

AZOPT contains benzalkonium chloride

AZOPT contains a preservative (benzalkonium chloride) which may cause eye irritation and is known to discolour soft contact lenses. Contact with soft contact lenses should be avoided. If you wear contact lenses you should remove them prior to the application of AZOPT and wait at least 15 minutes after instillation of the dose before putting your lenses back in.

3 How to use AZOPT

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.




Only use AZOPT for your eyes. Do not swallow or inject.

The recommended dose is

1 drop in the affected eye or eyes twice a day - morning and night.

Use this much unless your doctor told you to do something different. Only use AZOPT in both eyes if your doctor told you to. Take it for as long as your doctor told you to.

How to use



- Get the AZOPT bottle and a mirror
- Wash your hands
- Shake the bottle and twist off the cap. After the cap is removed, if the tamper evident snap collar is loose, remove before using product.
- Hold the bottle, pointing down, between your thumb and middle finger
- Tilt your head back. Pull down your eyelid with a clean finger, until there is a 'pocket' between the eyelid and your eye. The drop will go in here (picture 1)
- Bring the bottle tip close to the eye. Use the mirror if it helps
- Do not touch your eye or eyelid, surrounding areas or other surfaces with the dropper. It could infect the drops
- Gently press on the base of the bottle to release one drop of AZOPT at a time.
- Do not squeeze the bottle: it is designed so that a gentle press on the bottom is all that it needs (picture 2)
- After using AZOPT, press a finger to the corner of your eye, by the nose (picture 3) for at least 1 minute. This helps to stop AZOPT getting into the rest of the body.
- If you take drops in both eyes, repeat the steps for your other eye.
- Put the bottle cap back on firmly immediately after use
- Use up one bottle before opening the next bottle.

If a drop misses your eye, try again.

If you are using other eye drops, leave at least 5 minutes between putting in AZOPT and the other drops. Eye ointments should be administered last.

If you use more AZOPT than you should

If you get too much in your eyes, rinse it all out with warm water. Do not put in any more drops until it's time for your next regular dose.

If you forget to use AZOPT

Use a single drop as soon as you remember, and then go back to your regular routine. Do not use a double dose to make up for a forgotten dose.

If you stop using AZOPT

If you stop using AZOPT without speaking to your doctor, the pressure in your eye will not be controlled which could lead to loss of sight.

Turn the page for more advice

Now turn over ►

4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everyone gets them.

The following side effects have been seen with AZOPT®.

Common side effects

(may affect up to 1 in 10 people)

- **Effects in the eye:** blurred vision, eye irritation, eye pain, eye discharge, itchy eye, dry eye, abnormal eye sensation, redness of the eye.
- **General side effects:** bad taste.

Uncommon side effects

(may affect up to 1 in 100 people)

- **Effects in the eye:** sensitivity to light, inflammation or infection of the conjunctiva, eye swelling, eyelid itching, redness or swelling, growth on surface of eye, increased pigmentation of the eye, tired eyes, eyelid crusting, or increased tear production.
- **General side effects:** decreased or reduced heart function, palpitations, decreased heart rate, difficulty breathing, shortness of breath, cough, decreased red blood cell count in blood, increased chlorine level in blood, dizziness, drowsiness, difficulty with memory, depression, nervousness, generalized weakness, fatigue, feeling abnormal, pain, shaking, decreased sex drive, male sexual difficulty, cold symptoms, chest congestion, sinus infection, throat irritation, throat pain, abnormal or decreased sensation in mouth, inflammation of the lining of the oesophagus, abdominal pain, nausea, vomiting, upset stomach, frequent bowel movements, diarrhoea, intestinal gas, digestive disorder, kidney pain, muscle pain, muscle spasms, back pain, nose bleeds, runny nose, stuffy nose, sneezing, rash, abnormal skin sensation, itching, headache, dry mouth.

Rare side effects

(may affect up to 1 in 1,000 people)

- **Effects in the eye:** corneal swelling, double or reduced vision, abnormal vision, decreased eye sensation, swelling around the eye, increased pressure in eye, damage to the optic nerve.

- **General side effects:** memory impairment, drowsiness, chest pain, upper respiratory tract congestion, sinus congestion, nasal congestion, dry nose, ringing in ears, hair loss, generalized itching, feeling jittery, irritability, irregular heart rate, body weakness, difficulty sleeping.

Not known (frequency cannot be estimated from the available data):

- **Effects in the eye:** eyelid abnormality, visual disturbance, corneal disorder, eye allergy, decreased growth or number of eyelashes.
- **General side effects:** increased allergic symptoms, decreased sensation, tremor, loss or decrease in taste, decreased blood pressure, increased blood pressure, increased heart rate, joint pain, asthma, pain in extremity, skin redness, inflammation, or itching, abnormal liver blood tests, swelling of the extremities, frequent urination, decreased appetite.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed below. By reporting side effects you can help provide more information on the safety of this medicine.

Ireland

HPRA Pharmacovigilance,
Earlsfort Terrace, IRL - Dublin 2;
Tel: +353 1 6764971
Fax: +353 1 6762517.
Website: www.hpra.ie;
e-mail: medsafety@hpra.ie

Malta

ADR Reporting
The Medicines Authority
Post-Licensing Directorate
203 Level 3, Rue D'Argens
GŻR-1368 Gżira
Website: www.medicinesauthority.gov.mt
e-mail: postlicensing.medicinesauthority@gov.mt

United Kingdom

Yellow Card Scheme
Website: www.mhra.gov.uk/yellowcard

5 How to store AZOPT

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle and box after "EXP". The expiry date refers to the last day of the month.

This medicine does not require any special storage conditions.

You must throw away a bottle four weeks after you first opened it, to prevent infections. Write down the date you opened each bottle in the space below and in the space on the bottle label and box. For a pack containing a single bottle, write only one date.

Opened (1):
Opened (2):
Opened (3):

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 Contents of the pack and other information

What AZOPT contains

- The active substance is brinzolamide. Each milliliter contains 10 mg of brinzolamide.
- The other ingredients are benzalkonium chloride, carbomer 974P, edetate disodium, mannitol (E421), purified water, sodium chloride, tyloxapol. Tiny amounts of hydrochloric acid or sodium hydroxide are added to keep acidity levels (pH levels) normal.

What AZOPT looks like and contents of the pack

AZOPT is a milky liquid (a suspension) supplied in a pack containing a 5 ml or a 10 ml plastic (droptainer) bottle with a screw cap, or in a pack containing three 5 ml plastic (droptainer) bottles with screw caps. Not all pack sizes may be marketed.

Marketing Authorisation Holder

Alcon Laboratories (UK) Ltd., Frimley Business Park, Frimley, Camberley, Surrey GU16 7SR, United Kingdom

Manufacturer

S.A. Alcon-Couvreur N.V., Rijksweg 14, B-2870 Puurs, Belgium

Alcon[®]
a Novartis company

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien
Luxembourg/Luxemburg
Alcon NV
☎ +32 (0)2 754 32 10
(België/Belgique/Belgien)

България
Алкон България ЕООД
☎ +359 2 950 15 65

Česká republika
Alcon Pharmaceuticals
(Czech Republic) s.r.o.
☎ +420 225 775 111

Danmark
Alcon Nordic A/S
☎ +45 3636 4300

Deutschland
Alcon Pharma GmbH
☎ +49 (0)761 1304-0

Ελλάδα/Κύπρος
Άλκον Αις/Κυπριακή
Ελλάς ΑΕΒΕ
☎ +30 210 68 78 300
(Ελλάδα)

Eesti
Alcon Pharmaceuticals Ltd.
Eesti filiaal
☎ +372 6 313 214

España
Alcon Cusi, S.A.
☎ +34 93 497 7000

France
Laboratoires Alcon
☎ +33 (0)1 47 10 47 10

Hrvatska
Alcon Farmaceutika d.o.o.
☎ +385 1 4611 988

Ireland
Malta
United Kingdom
Alcon Laboratories (UK) Ltd
☎ +44 (0)871 376 1402
(United Kingdom)

Ísland
Alcon Nordic A/S
☎ +45 3636 4300

Italia
Alcon Italia S.p.A.
☎ +39 02 81 80 31

Latvija
Alcon Pharmaceuticals Ltd
☎ +371 67 321 121

Lietuva
Alcon Pharmaceuticals Ltd
atstovybė
☎ +370 5 2 314 756

Magyarország
Alcon Hungária
Gyógyszerkereskedelmi Kft
☎ +36-1-463-9080

Nederland
Alcon Nederland BV
☎ +31 (0) 183 654321

Norge
Alcon Nordic A/S
☎ +45 3636 4300

Österreich
Alcon Ophthalmika GmbH
☎ +43 (0)1 596 69 70

Polska
Alcon Polska Sp. z o.o.
☎ +48 22 820 3450

Portugal
Alcon Portugal -
Produtos e Equipamentos
Oftalmológicos, Lda.
☎ +351 214 400 300

România
S.C. Alcon Romania S.R.L.
☎ +40 21 203 93 24

Slovenija
Alcon d.o.o.
☎ +386 1 422 5280

Slovenská republika
Novartis Slovakia s.r.o.
Alcon Division
☎ +421 2 5441 0378

Suomi/Finland
Alcon Nordic A/S
☎ +45 3636 4300

Sverige
Alcon Nordic A/S
☎ +45 3636 4300

This leaflet was last revised in April 2014

Detailed information on this product is available on the website of the European Medicines Agency (EMA) <http://www.ema.europa.eu>